

A Study on ca – Domination of Cartesian product of a Class of Path Semigraphs

N. Murugesan¹, D. Narmatha²

¹(Mathematics, Government Arts College, Coimbatore, India)

²(Mathematics, Sri Ramakrishna Engineering College, Coimbatore, India)

Abstract: In this paper, we study ca – domination number of the cartesian product of some simple path semigraphs.

Keywords: Cartesian Product, Domination number, ca – Domination number, Path Semigraph, Semigraph.

1. Introduction

A semigraph S is a pair (V, X) where V is a nonempty set whose elements are called vertices of S and X is a set of ordered n -tuples $n \geq 2$ of distinct vertices called edges of S satisfying the following conditions :

- i. any two edges have at most one vertex in common.
- ii. two edges $E_1 = (u_1, u_2, \dots, u_m)$ and $E_2 = (v_1, v_2, \dots, v_n)$ are said to be equal iff
 - a. $m = n$ and
 - b. either $u_i = v_i$ or $u_i = v_{n-i+1}$ for $1 \leq i \leq n$.

Semigraphs are introduced by E. Sampath kumar [8] in the year 2000, since then it has been an interesting field of research in graph theory. B. D. Acharya [1] discussed construction of semigraph from square matrices. B. Y. Bam and N. S. Bhavne [2] have studied different types of degree sequences in semigraphs. S. P. Subbiah [9] have studied the relationship between topologies and discrete semigraphs.

There are different types of adjacency defined between two vertices in a semigraph. Two vertices u and v in a semigraph S are said to be ca – adjacent if they are adjacent and consecutive in order as well. The basic concepts relating semigraphs are discussed in [4]. The properties of various types of adjacencies between vertices are deeply discussed in [5].

A subset D of V is said to be ca – dominating set if for every $v \in V - D$ there exists an vertex $u \in D$ such that u and v are consecutively adjacent. The minimum cardinality of such a set D is called ca – domination number of the semigraph S . It is denoted as $\gamma_{ca}(S)$.

A path semigraph $S = (V, X)$ is a semigraph with the following properties.

- i. it has no middle – end vertices.
- ii. it has exactly two end vertices each with edge degree one.
- iii. the edge degree of all other end vertices (if they exist) are exactly two.

Note that a path semigraph with no middle vertices is simply a path. The certain energies of path semigraphs have been studied in [6]. Characteristic polynomial of path semigraphs have been studied in [7]. In this paper, we discuss the ca – domination number of cartesian product of path semigraphs.

2. Cartesian Product of Path Semigraphs

Let $S_1 = (V_1, X_1)$ and $S_2 = (V_2, X_2)$ be two semigraphs. The Cartesian product of S_1 and S_2 denoted by $S_1 \square S_2$ is defined as follows :

Vertex set of $S_1 \square S_2$ is $V_1 \times V_2$ and the edge set is as given below:

For any vertex $u \in V_1$ and any edge $E = (v_1, v_2, \dots, v_r)$ in X_2 , $((u, v_1), (u, v_2), \dots, (u, v_r))$ is an edge in $S_1 \square S_2$. Also, for any edge $E = (u_1, u_2, \dots, u_s)$ in X_1 and for any vertex $v \in V_2$, $((u_1, v), (u_2, v), \dots, (u_s, v))$ is an edge in $S_1 \square S_2$.

It has been discussed that Cartesian product of two semigraphs is also a semigraph [3].

2.1. Example

Consider the following two semigraphs.




Fig. 2.1 two semigraphs S_1, S_2

The cartesian product of the above two semigraphs is given in fig. 2.2.




Fig. 2.2 cartesian product $S_1 \square S_2$

In this paper, we study the ca – domination number of the cartesian product of simple path semigraphs. A simple path semigraph means, the path semigraph in which every edge contains exactly one middle vertex. A simple path semigraph with n edges is represented by $P_{s(n)}$. In this paper, we study ca – domination number of $P_{s(n)} \square P_{s(m)}$, for $n = 1, 2, \dots$ and $m = 1, 2, 3, \dots, 7$.

2.2 Lemma

$$\begin{aligned}
 & \text{i. } \gamma_{ca} [P_{s(1)} \square P_{s(1)}] = 3 \\
 & \text{ii. } \gamma_{ca} [P_{s(n)} \square P_{s(1)}] = \begin{cases} 3 \binom{n-2}{2} + 4 & \text{if } n = 2k + 2 \\ 3 \binom{n-3}{2} + 6 & \text{if } n = 2k + 3 \\ k = 0, 1, 2, \dots \end{cases}
 \end{aligned}$$

Proof :

Let $P_{s(1)}$ represents the simple path semigraph having exactly one middle vertex. The cartesian product $P_{s(1)} \square P_{s(1)}$, is given in the following fig.2.3.



Fig. 2.3 $P_{s(1)}$

The vertex set and edge set of the cartesian product $P_{s(1)} \square P_{s(1)}$ is $V = \{(u_1, v_1), (u_2, v_1), (u_3, v_1), (u_1, v_2), (u_2, v_2), (u_3, v_2), (u_1, v_3), (u_2, v_3), (u_3, v_3)\}$

$$E = \left\{ \left[(u_1, v_1), (u_2, v_1), (u_3, v_1) \right], \left[(u_1, v_2), (u_2, v_2), (u_3, v_2) \right], \left[(u_1, v_3), (u_2, v_3), (u_3, v_3) \right], \right. \\ \left. \left[(u_1, v_1), (u_1, v_2), (u_1, v_3) \right], \left[(u_2, v_1), (u_2, v_2), (u_2, v_3) \right], \left[(u_3, v_1), (u_3, v_2), (u_3, v_3) \right] \right\}$$

In the fig. 2.2 (u_2, v_2) is the only middle vertex, and $(u_2, v_1), (u_1, v_2), (u_3, v_2), (u_2, v_3)$ are middle-end vertices, and $(u_1, v_1), (u_3, v_1), (u_1, v_3), (u_3, v_3)$ are end vertices.

Fig.2.4 shows the cartesian products of $P_{s(1)} \square P_{s(1)}$ and $P_{s(2)} \square P_{s(1)}$.




Fig. 2.4 $P_{s(1)} \square P_{s(1)} ; P_{s(2)} \square P_{s(1)}$

First let $k = 0$, the vertex set $\{(u_3, v_1), (u_3, v_3), (u_1, v_2)\}$ form a minimal ca – dominating set for $P_{s(1)} \square P_{s(1)}$. Hence, $\gamma_{ca} [P_{s(1)} \square P_{s(1)}] = 3$. This proves (i).

To prove (ii), let us take $n = 2k + 2, k = 0,1,2,..$

$$\text{The minimal } ca \text{ – dominating set is } D = \left\{ \begin{array}{l} \left\{ \bigcup_{l=1}^{k+1} (u_{2(2l-1)+1}, v_j) / j=1,3 \right\} \\ \bigcup_{l=0}^{k+1} (u_{4l+1}, v_2) \end{array} \right\}$$

Now

$$\begin{aligned} |D| &= 2(k+1) + k + 2 \\ &= 3k + 4 \\ &= 3 \left(\frac{n-2}{2} \right) + 4 \end{aligned}$$

Let $n = 2k + 3; k = 0,1,2,3,..$. The minimal ca – dominating set is

$$D = \left\{ \begin{array}{l} \left\{ \bigcup_{l=1}^{k+2} (u_{2(2l-1)+1}, v_j) / j=1,3 \right\} \\ \bigcup_{l=0}^{k+1} (u_{4l+1}, v_2) \end{array} \right\}$$

Hence

$$\begin{aligned} |D| &= 2(k+2) + k + 2 \\ &= 3k + 6 \\ &= 3 \left(\frac{n-3}{2} \right) + 6 \end{aligned}$$

Hence the theorem.

2.3. Theorem

i. $\gamma_{ca} [P_{s(1)} \square P_{s(2)}] = 4$

ii. $\gamma_{ca} [P_{s(2)} \square P_{s(2)}] = 7$

$$\text{iii. } \gamma_{ca} [P_{s(n)} \square P_{s(2)}] = \begin{cases} 5\left(\frac{n-3}{2}\right)+10 & \text{if } n = 2k + 3 \\ 5\left(\frac{n-4}{2}\right)+12 & \text{if } n = 2k + 4 \\ k = 0,1,2,\dots \end{cases}$$

Proof :

The vertex set $\{(u_2, v_1), (u_2, v_5), (u_1, v_3), (u_3, v_3)\}$ and $\{(u_3, v_1), (u_3, v_3), (u_3, v_5), (u_1, v_2), (u_5, v_2), (u_1, v_4), (u_5, v_4)\}$ are the minimal ca – dominating sets for $P_{s(1)} \square P_{s(2)}$, $P_{s(2)} \square P_{s(2)}$ respectively. This proves (i) and (ii).

To prove (iii), let us take $n = 2k + 3, k = 0,1,2,\dots$

$$\text{The minimal } ca \text{ – dominating set is } D = \left\{ \begin{array}{l} \left\{ \bigcup_{l=1}^{k+2} (u_{2(2l-1)+1}, v_j) / j=1,3,5 \right\} \\ \bigcup_{l=0}^{k+1} (u_{4l+1}, v_j) / j = 2,4. \end{array} \right\}$$

Now

$$\begin{aligned} |D| &= 3(k+2) + 2(k+2) \\ &= 5k + 10 \\ &= 5\left(\frac{n-3}{2}\right) + 10 \end{aligned}$$

Let $n = 2k + 4; k = 0,1,2,3,\dots$. The minimal ca – dominating set is

$$D = \left\{ \begin{array}{l} \left\{ \bigcup_{l=1}^{k+2} (u_{2(2l-1)+1}, v_j) / j=1,3,5 \right\} \\ \bigcup_{l=0}^{k+2} (u_{4l+1}, v_j) / j = 2,4 \end{array} \right\}$$

Hence

$$\begin{aligned} |D| &= 3(k+2) + 2(k+3) \\ &= 5k + 12 \\ &= 5\left(\frac{n-4}{2}\right) + 12 \end{aligned}$$

Hence the theorem.

2.4 Theorem

i. $\gamma_{ca} [P_{s(1)} \square P_{s(3)}] = 6$

ii. $\gamma_{ca} [P_{s(2)} \square P_{s(3)}] = 10$

iii. $\gamma_{ca} [P_{s(3)} \square P_{s(3)}] = 14$

$$iv. \gamma_{ca} [P_{s(n)} \square P_{s(3)}] = \begin{cases} 7\left(\frac{n-4}{2}\right) + 17 & \text{if } n = 2k + 4 \\ 7\left(\frac{n-5}{2}\right) + 21 & \text{if } n = 2k + 5 \\ k = 0,1,2,\dots \end{cases}$$

Proof:

The vertex set $\{(u_2, v_1), (u_2, v_5), (u_1, v_3), (u_3, v_3), (u_1, v_7), (u_3, v_7)\}$ and $\{(u_2, v_1), (u_4, v_1), (u_1, v_3), (u_3, v_3), (u_5, v_3), (u_2, v_5), (u_4, v_5), (u_1, v_7), (u_3, v_7), (u_5, v_7)\}$ are minimal *ca* – dominating sets for $P_{s(1)} \square P_{s(3)}$, $P_{s(2)} \square P_{s(3)}$ respectively. This proves (i) and (ii).

The vertex set

$$\left\{ \begin{array}{l} (u_3, v_1), (u_7, v_1), (u_1, v_2), (u_5, v_2), (u_3, v_3), (u_7, v_3), (u_1, v_4), (u_5, v_4), (u_3, v_5), (u_7, v_5), \\ (u_1, v_6), (u_5, v_6), (u_3, v_7), (u_7, v_7) \end{array} \right\}$$

is a minimal *ca* – dominating sets for $P_{s(3)} \square P_{s(3)}$. This proves (iii).

To prove (iv), let us take $n = 2k + 4, k = 0,1,2,\dots$. Here

$$D = \left\{ \begin{array}{l} \left\{ \bigcup_{l=1}^{k+2} (u_{2(2l-1)+1}, v_j) / j = 1,3,5,7 \right\} \\ \bigcup_{l=0}^{k+2} (u_{4l+1}, v_j) / j = 2,4,6. \end{array} \right\}$$

Now

$$\begin{aligned} |D| &= 3(k+3) + 4(k+2) \\ &= 7k + 17 \\ &= 7\left(\frac{n-4}{2}\right) + 17 \end{aligned}$$

Let $n = 2k + 5; k = 0,1,2,3,\dots$. Here

$$D = \left\{ \begin{array}{l} \left\{ \bigcup_{l=1}^{k+3} (u_{2(2l-1)+1}, v_j) / j = 1,3,5,7 \right\} \\ \bigcup_{l=0}^{k+2} (u_{4l+1}, v_j) / j = 2,4,6 \end{array} \right\}$$

Hence

$$\begin{aligned} |D| &= 3(k+3) + 4(k+3) \\ &= 7k + 21 \\ &= 7\left(\frac{n-5}{2}\right) + 21 \end{aligned}$$

Hence the theorem.

2.5 Theorem

i. $\gamma_{ca} [P_{s(1)} \square P_{s(4)}] = 7$

ii. $\gamma_{ca} [P_{s(2)} \square P_{s(4)}] = 12$

iii. $\gamma_{ca} [P_{s(3)} \square P_{s(4)}] = 17$

iv. $\gamma_{ca} [P_{s(4)} \square P_{s(4)}] = 22$

$$v. \gamma_{ca} [P_{s(n)} \square P_{s(4)}] = \begin{cases} 9\left(\frac{n-5}{2}\right) + 27 & \text{if } n = 2k + 5 \\ 9\left(\frac{n-6}{2}\right) + 31 & \text{if } n = 2k + 6 \\ k = 0,1,2,\dots \end{cases}$$

Proof:

The vertex set $\{(u_2, v_1), (u_2, v_5), (u_1, v_3), (u_3, v_3), (u_1, v_7), (u_3, v_7), (u_2, v_9)\}$ form a minimal *ca* – dominating sets for $P_{s(1)} \square P_{s(4)}$ respectively. This proves (i).

The vertex set $\{(u_i, v_j) / i = 2,4, j = 1,5,9\} \& \{(u_i, v_j) / i = 1,3,5, j = 3,7\}$ is a minimal *ca* – dominating sets for $P_{s(2)} \square P_{s(4)}$. This proves (ii).

The vertex set $\{(u_i, v_j) / i = 2,4,6, j = 1,5,9\} \& \{(u_i, v_j) / i = 1,3,5,7, j = 3,7\}$ is a minimal *ca* – dominating sets for $P_{s(3)} \square P_{s(4)}$. This proves (iii).

The vertex set $\{(u_i, v_j) / i = 2,4,6,8, j = 1,5,9\} \& \{(u_i, v_j) / i = 1,3,5,9, j = 3,7\}$

is a minimal *ca* – dominating sets for $P_{s(4)} \square P_{s(4)}$. This proves (iv).

To prove (iv), let us take $n = 2k + 5, k = 0,1,2,\dots$ Here

$$D = \left\{ \begin{array}{l} \left\{ \bigcup_{l=1}^{k+3} (u_{2(2l-1)+1}, v_j) / j = 1,3,5,7,9 \right\} \\ \bigcup_{l=0}^{k+3} (u_{4l+1}, v_j) / j = 2,4,6,8. \end{array} \right\}$$

Now

$$\begin{aligned} |D| &= 4(k+3) + 5(k+3) \\ &= 9k + 27 \\ &= 9\left(\frac{n-5}{2}\right) + 27 \end{aligned}$$

Let $n = 2k + 6; k = 0,1,2,3,\dots$ Here

$$D = \left\{ \begin{array}{l} \left\{ \bigcup_{l=1}^{k+3} (u_{2(2l-1)+1}, v_j) / j = 1,3,5,7,9 \right\} \\ \bigcup_{l=0}^{k+3} (u_{4l+1}, v_j) / j = 2,4,6,8 \end{array} \right\}$$

Hence

$$\begin{aligned} |D| &= 4(k+4) + 5(k+3) \\ &= 9k + 31 \\ &= 9\left(\frac{n-6}{2}\right) + 31 \end{aligned}$$

Hence the theorem.

2.6 Theorem

- i. $\gamma_{ca} [P_{s(1)} \square P_{s(5)}] = 9$
- ii. $\gamma_{ca} [P_{s(2)} \square P_{s(5)}] = 15$
- iii. $\gamma_{ca} [P_{s(3)} \square P_{s(5)}] = 21$

- iv. $\gamma_{ca} [P_{s(4)} \square P_{s(5)}] = 27$
- v. $\gamma_{ca} [P_{s(5)} \square P_{s(5)}] = 33$

- vi. $\gamma_{ca} [P_{s(n)} \square P_{s(5)}] = \begin{cases} 11 \left(\frac{n-6}{2} \right) + 38 & \text{if } n = 2k + 6 \\ 11 \left(\frac{n-7}{2} \right) + 44 & \text{if } n = 2k + 7 \\ k = 0,1,2,\dots \end{cases}$

Proof:

The vertex set $\{ (u_2, v_j) / j = 1,5,9 \} \& \{ (u_i, v_j) / i = 1,3, j = 3,7,11 \}$ form a minimal $ca -$ dominating sets for $P_{s(1)} \square P_{s(5)}$ respectively. This proves (i).

The vertex set $\{ (u_i, v_j) / i = 2,4, j = 1,5,9 \} \& \{ (u_i, v_j) / i = 1,3,5, j = 3,7,11 \}$ is a a minimal $ca -$ dominating sets for $P_{s(2)} \square P_{s(5)}$. This proves (ii).

The vertex set $\{ (u_i, v_j) / i = 2,4,6, j = 1,5,9 \} \& \{ (u_i, v_j) / i = 1,3,5,7, j = 3,7,11 \}$ is a a minimal $ca -$ dominating sets for $P_{s(3)} \square P_{s(5)}$. This proves (iii).

The vertex set $\{ (u_i, v_j) / i = 2,4,6,8, j = 1,5,9 \} \& \{ (u_i, v_j) / i = 1,3,5,7,9, j = 3,7,11 \}$ is a a minimal $ca -$ dominating sets for $P_{s(4)} \square P_{s(5)}$. This proves (iv).

The vertex set $\{ (u_i, v_j) / i = 2,4,6,8,10, j = 1,5,9 \} \& \{ (u_i, v_j) / i = 1,3,5,7,9,11, j = 3,7,11 \}$ is a a minimal $ca -$ dominating sets for $P_{s(5)} \square P_{s(5)}$. This proves (v).

To prove (vi), let us take $n = 2k + 6, k = 0,1,2,..$ Here

$$D = \left\{ \begin{array}{l} \left\{ \bigcup_{l=1}^{k+3} (u_{2(2l-1)+1}, v_j) / j = 1,3,5,7,9,11 \right\} \\ \left\{ \bigcup_{l=0}^{k+3} (u_{4l+1}, v_j) / j = 2,4,6,8,10. \right\} \end{array} \right\}$$

Now

$$\begin{aligned} |D| &= 5(k+4) + 6(k+3) \\ &= 11k + 38 \\ &= 11 \left(\frac{n-6}{2} \right) + 38 \end{aligned}$$

Let $n = 2k + 7; k = 0,1,2,3,\dots$. Here

$$D = \left\{ \begin{array}{l} \left\{ \bigcup_{l=1}^{k+4} (u_{2(2l-1)+1}, v_j) / j=1,3,5,7,9,11 \right\} \\ \left\{ \bigcup_{l=0}^{k+3} (u_{4l+1}, v_j) / j=2,4,6,8,10. \right\} \end{array} \right\}$$

Hence

$$\begin{aligned} |D| &= 5(k+4) + 6(k+4) \\ &= 11k + 44 \\ &= 11 \left(\frac{n-7}{2} \right) + 44 \end{aligned}$$

Hence the theorem.

2.7 Theorem

- i. $\gamma_{ca} [P_{s(1)} \square P_{s(6)}] = 10$
- ii. $\gamma_{ca} [P_{s(2)} \square P_{s(6)}] = 17$
- iii. $\gamma_{ca} [P_{s(3)} \square P_{s(6)}] = 24$
- iv. $\gamma_{ca} [P_{s(4)} \square P_{s(6)}] = 31$
- v. $\gamma_{ca} [P_{s(5)} \square P_{s(6)}] = 38$
- vi. $\gamma_{ca} [P_{s(6)} \square P_{s(6)}] = 45$
- vii. $\gamma_{ca} [P_{s(n)} \square P_{s(6)}] = \begin{cases} 13 \left(\frac{n-7}{2} \right) + 52 & \text{if } n = 2k + 7 \\ 13 \left(\frac{n-8}{2} \right) + 58 & \text{if } n = 2k + 8 \\ k = 0,1,2,\dots \end{cases}$

Proof:

The vertex set $\{(u_2, v_j) / j = 1,5,9,13\} \& \{(u_i, v_j) / i = 1,3, j = 3,7,11\}$ form a minimal *ca* – dominating sets for $P_{s(1)} \square P_{s(6)}$ respectively. This proves (i).

The vertex set $\{(u_i, v_j) / i = 2,4, j = 1,5,9,13\} \& \{(u_i, v_j) / i = 1,3,5, j = 3,7,11\}$ is a a minimal *ca* – dominating sets for $P_{s(2)} \square P_{s(6)}$. This proves (ii).

The vertex set $\{(u_i, v_j) / i = 2,4,6, j = 1,5,9,13\} \& \{(u_i, v_j) / i = 1,3,5,7, j = 3,7,11\}$ is a a minimal *ca* – dominating sets for $P_{s(3)} \square P_{s(6)}$. This proves (iii).

The vertex set $\{(u_i, v_j) / i = 2,4,6,8, j = 1,5,9,13\} \& \{(u_i, v_j) / i = 1,3,5,7,9, j = 3,7,11\}$ is a a minimal *ca* – dominating sets for $P_{s(4)} \square P_{s(6)}$. This proves (iv).

The vertex set $\{(u_i, v_j) / i = 2,4,6,8,10, j = 1,5,9,13\} \& \{(u_i, v_j) / i = 1,3,5,7,9,11, j = 3,7,11\}$ is a a minimal *ca* – dominating sets for $P_{s(5)} \square P_{s(6)}$. This proves (v).

The vertex set $\{(u_i, v_j) / i = 2,4,6,8,10,12, j = 1,5,9,13\} \& \{(u_i, v_j) / i = 1,3,5,7,9,11,13, j = 3,7,11\}$ is a a minimal *ca* – dominating sets for $P_{s(6)} \square P_{s(6)}$. This proves (vi).

To prove (vii), let us take $n = 2k + 7, k = 0,1,2,\dots$ Here

$$D = \left\{ \begin{array}{l} \left\{ \bigcup_{l=1}^{k+7} (u_{2l}, v_j) / j=1,5,13,19 \right\} \\ \left\{ \bigcup_{l=0}^{k+7} (u_{2l+1}, v_j) / j = 3,7,11. \right\} \end{array} \right\}$$

Now

$$\begin{aligned} |D| &= 8(k+3) + 4(k+7) \\ &= 13k + 52 \\ &= 13 \left(\frac{n-7}{2} \right) + 52 \end{aligned}$$

Let $n = 2k + 8; k = 0,1,2,3, \dots$. Here

$$D = \left\{ \begin{array}{l} \left\{ \bigcup_{l=1}^{k+4} (u_{2(2l-1)+1}, v_j) / j=1,3,5,7,9,11,13 \right\} \\ \left\{ \bigcup_{l=0}^{k+4} (u_{4l+1}, v_j) / j = 2,5,9, \dots, (4k+1). \right\} \end{array} \right\}$$

Hence

$$\begin{aligned} |D| &= 6(k+5) + 7(k+4) \\ &= 13k + 58 \\ &= 13 \left(\frac{n-8}{2} \right) + 58 \end{aligned}$$

Hence the theorem.

2.8 Theorem

- i. $\gamma_{ca} [P_{s(1)} \square P_{s(7)}] = 12$
- ii. $\gamma_{ca} [P_{s(2)} \square P_{s(7)}] = 20$
- iii. $\gamma_{ca} [P_{s(3)} \square P_{s(7)}] = 28$
- iv. $\gamma_{ca} [P_{s(4)} \square P_{s(7)}] = 36$
- v. $\gamma_{ca} [P_{s(5)} \square P_{s(7)}] = 44$
- vi. $\gamma_{ca} [P_{s(6)} \square P_{s(7)}] = 52$
- vii. $\gamma_{ca} [P_{s(7)} \square P_{s(7)}] = 60$

$$\text{viii. } \gamma_{ca} [P_{s(n)} \square P_{s(7)}] = \begin{cases} 15 \left(\frac{n-8}{2} \right) + 67 & \text{if } n = 2k + 8 \\ 15 \left(\frac{n-9}{2} \right) + 75 & \text{if } n = 2k + 9 \\ k = 0,1,2, \dots \end{cases}$$

Proof:

The vertex set $\{(u_2, v_j) / j = 1,5,9,13\} \& \{(u_i, v_j) / i = 1,3, j = 3,7,11,15\}$ form a minimal ca – dominating sets for $P_{s(1)} \square P_{s(7)}$ respectively. This proves (i).

The vertex set $\{(u_i, v_j) / i = 2, 4, j = 1, 5, 9, 13\} \& \{(u_i, v_j) / i = 1, 3, 5, j = 3, 7, 11, 15\}$

is a a minimal ca – dominating sets for $P_{s(2)} \square P_{s(7)}$. This proves (ii).

The vertex set $\{(u_i, v_j) / i = 2, 4, 6, j = 1, 5, 9, 13\} \& \{(u_i, v_j) / i = 1, 3, 5, 7, j = 3, 7, 11, 15\}$

is a a minimal ca – dominating sets for $P_{s(3)} \square P_{s(7)}$. This proves (iii).

The vertex set $\{(u_i, v_j) / i = 2, 4, 6, 8, j = 1, 5, 9, 13\} \& \{(u_i, v_j) / i = 1, 3, 5, 7, 9, j = 3, 7, 11, 15\}$

is a a minimal ca – dominating sets for $P_{s(4)} \square P_{s(7)}$. This proves (iv).

The vertex set

$\{(u_i, v_j) / i = 2, 4, 6, 8, 10, j = 1, 5, 9, 13\} \& \{(u_i, v_j) / i = 1, 3, 5, 7, 9, 11, j = 3, 7, 11, 15\}$

is a a minimal ca – dominating sets for $P_{s(5)} \square P_{s(7)}$. This proves (v).

The vertex set

$\{(u_i, v_j) / i = 2, 4, 6, 8, 10, 12, j = 1, 5, 9, 13\} \& \{(u_i, v_j) / i = 1, 3, 5, 7, 9, 11, 13, j = 3, 7, 11, 15\}$

is a a minimal ca – dominating sets for $P_{s(6)} \square P_{s(7)}$. This proves (vi).

The vertex set

$\{(u_i, v_j) / i = 2, 4, 6, 8, 10, 12, j = 1, 5, 9, 13\} \& \{(u_i, v_j) / i = 1, 3, 5, 7, 9, 11, 13, 15, j = 3, 7, 11, 15\}$

is a a minimal ca – dominating sets for $P_{s(7)} \square P_{s(7)}$. This proves (vii).

To prove (viii), let us take $n = 2k + 8, k = 0, 1, 2, \dots$. Here

$$D = \left\{ \begin{array}{l} \left\{ \bigcup_{l=1}^{k+4} (u_{2(2l-1)+1}, v_j) / j = 1, 3, 5, 7, 9, 11, 13, 15 \right\} \\ \bigcup_{l=0}^{k+4} (u_{4l+1}, v_j) / j = 2, 4, 6, 8, 10, 12, 14. \end{array} \right\}$$

Now

$$\begin{aligned} |D| &= 8(k+4) + 7(k+5) \\ &= 15k + 67 \\ &= 15 \left(\frac{n-8}{2} \right) + 67 \end{aligned}$$

Let $n = 2k + 9; k = 0, 1, 2, 3, \dots$. Here

$$D = \left\{ \begin{array}{l} \left\{ \bigcup_{l=1}^{k+5} (u_{2(2l-1)+1}, v_j) / j = 1, 3, 5, 7, 9, 11, 13, 15 \right\} \\ \bigcup_{l=0}^{k+4} (u_{4l+1}, v_j) / j = 2, 4, 6, 8, 10, 12, 14. \end{array} \right\}$$

Hence

$$\begin{aligned} |D| &= 8(k+5) + 7(k+5) \\ &= 15k + 75 \\ &= 15 \left(\frac{n-9}{2} \right) + 75 \end{aligned}$$

Hence the theorem.

3. Conclusion

It has been interesting fact to study ca – domination number of $P_{s(n)} \square P_{s(m)}$.

In this paper, it has been discussed the ca – domination number of cartesian product graphs $P_{s(n)} \square P_{s(m)}$, $m=1,2,\dots,7$ but for all values of n .

References

- [1]. E. Sampathkumar, Semigraphs and their Applications, *Report on the DST project*, submitted to DST, India, May 2000.
- [2]. B. D. Acharya, Semigraphic Matrices and Matrix Semigraph, Private Communication.
- [3]. B. Y. Bam and N. S. Bhav, *On Some Problems of Graph Theory in Semigraphs*, Ph.D Thesis, University of Pune.
- [4]. S. P. Subbaih, *A Study of Graph Theory: Topology, Steiner Domination and Semigraph Concepts*, Ph.D Thesis, Madurai Kamaraj University, 2007.
- [5]. F. Harary, *Graph Theory*, Addition Wesley, Reading Mass, 1972.
- [6]. N. Murugesan and D. Narmatha, Some properties of Semigraph and its Associated Graphs, *International Journal of Engineering Research and Technology*, Vol. 3(5), 2014, 898-903.
- [7]. N. Murugesan and D. Narmatha, e and n – Energies of Linear Semigraph, *International Journal of Research in Engineering and Technology*, Vol. 3(11), 2014, 174-178.
- [8]. N. Murugesan and D. Narmatha, Characteristic Polynomial of r -adjacency matrix of Path Semigraph, *International Journal of Innovative Research in Science, Engineering and Technology*, Vol. 5(7), 2016, 1111-1119.
- [9]. S. Gomathi, *Studies in Semigraphs and Domination*, Ph.D Thesis, Madurai Kamaraj University, 2008.